

(EMC01-0001E)


- If $(x - a)$ is a factor of the polynomial $x^6 - ax^5 + x^4 - ax^3 + 3x + a - 2$, then the value of a is
 - A) 1
 - B) $\frac{1}{2}$
 - C) $\frac{1}{3}$
 - D) d)2

(EMC01-0002M)

- The geometric mean of the roots of the equation $x^2 - 18x + 9 = 0$ is
 - A) 3
 - B) $3\sqrt{2}$
 - C) 9
 - D) $9\sqrt{2}$

(EMC01-0003M)

- If the roots of $x^2 + ax + 9 = 0$ are complex, then
 - A) $a < -6$
 - B) $a < 6$
 - C) $|a| < 6$
 - D) $|a| > 6$



(EMC01-0004M)

- The value of k for which the quadratic equation $x^2 - 2x(1 + 3k) + 7(2k + 3) = 0$ has equal roots, is
 - A) 1
 - B) 2
 - C) 3
 - D) 4

(EMC01-0005E)

- If the product of roots of the equation $mx^2 + 6x + (2m - 1) = 0$ is -1 , then the value of m is
 - A) -1
 - B) $-\frac{1}{3}$
 - C) $\frac{1}{3}$
 - D) 1

(EMC01-0006E)

- If a and b are roots of $x^2 - px + q = 0$, then $\frac{1}{a} + \frac{1}{b}$ is equal to
 - A) $\frac{-p}{q}$
 - B) $\frac{1}{2p}$
 - C) $\frac{-1}{q}$
 - D) $\frac{p}{q}$

(EMC01-0007M)

- One root of the equation $5x^2 + 13x + m = 0$ is reciprocal of the other if m equals
 - A) 0
 - B) 5
 - C) $\frac{1}{6}$
 - D) 6

(EMC01-0008M)

- If $1 - i$ is a root of the equation $x^2 + ax + b = 0$, then b is equal to
 - A) -2
 - B) -1
 - C) 1
 - D) 2

(EMC01-0010H)

- If α and β are the roots of the equation $x^2 + 2x + 4 = 0$, then $\frac{1}{\alpha^3} + \frac{1}{\beta^3}$ is equal to

- A) $-\frac{1}{2}$
- B) $\frac{1}{2}$
- C) $\frac{1}{4}$
- D) $\frac{1}{6}$

(EMC01-0011H)

- Let $p, q \in \{1, 2, 3, 4, 5\}$. The number of equations of the form $px^2 + qx + 1 = 0$ having real roots, is

- A) 7
- B) 8
- C) 9
- D) 12

(EMC01-0012M)

- If the difference of the roots of the equation $x^2 + px + 8 = 0$ is 2, then p equals

- A) ± 2
- B) $-6, 2$
- C) $-2, 6$
- D) ± 6

(EMC01-0013M)

- If one root of the equation $x^2 + px + q = 0$ is $2 + \sqrt{3}$, then the values of p and q are

- A) $-2, -\sqrt{3}$
- B) $-4, 1$
- C) $2, \sqrt{3}$
- D) $4, -1$

(EMC01-0014M)

- For what value of m , the ratio of the roots of the equation $12x^2 - mx + 5 = 0$ is 3 : 2?

A) $5\sqrt{10}$
B) $10\sqrt{5}$
C) $25\sqrt{2}$
D) $15\sqrt{5}$

(EMC01-0016H)


- The roots of equation $2^{2x} - 10 \cdot 2^x + 16 = 0$ are

A) 1, 3
B) 1, 8
C) 2, 3
D) 2, 8

(EMC01-0017M)

- The solution set of $\sqrt{2x - 6} + \sqrt{x + 4} = 5$ is

A) {5}
B) {0, 5}
C) {1, 3}
D) {3, 5}



PARHO AUR
Chaa�ao

Answers Key	
1	B
2	A
3	C
4	B
5	C
6	D
7	B
8	D
10	C
11	D
12	D
13	B
14	A
16	A
17	A