

(EMC01-0001E)

- If $(x - a)$ is a factor of the polynomial $x^6 - ax^5 + x^4 - ax^3 + 3x + a - 2$, then the value of a is
 - A) 1
 - B) $\frac{1}{2}$
 - C) $\frac{1}{3}$
 - D) 2

(EMC01-0002M)

- The geometric mean of the roots of the equation $x^2 - 18x + 9 = 0$ is
 - A) 3
 - B) $3\sqrt{2}$
 - C) 9
 - D) $9\sqrt{2}$

(EMC01-0003M)

- If the roots of $x^2 + ax + 9 = 0$ are complex, then
 - A) $a < -6$
 - B) $a < 6$
 - C) $|a| < 6$
 - D) $|a| > 6$

(EMC01-0004M)

- The value of k for which the quadratic equation $x^2 - 2x(1 + 3k) + 7(2k + 3) = 0$ has equal roots, is
 - A) 1
 - B) 2
 - C) 3
 - D) 4

(EMC01-0005E)

- If the product of roots of the equation $mx^2 + 6x + (2m - 1) = 0$ is -1 , then the value of m is
 - A) -1
 - B) $-\frac{1}{3}$
 - C) $\frac{1}{3}$
 - D) 1

(EMC01-0006E)

- If a and b are roots of $x^2 - px + q = 0$, then $\frac{1}{a} + \frac{1}{b}$ is equal to
 - A) $\frac{-p}{q}$
 - B) $\frac{1}{2p}$
 - C) $\frac{-1}{q}$
 - D) $\frac{p}{q}$

(EMC01-0007M)

- One root of the equation $5x^2 + 13x + m = 0$ is reciprocal of the other if m equals
 - A) 0
 - B) 5
 - C) $\frac{1}{6}$
 - D) 6

(EMC01-0008M)

- If $1 - i$ is a root of the equation $x^2 + ax + b = 0$, then b is equal to
 - A) -2
 - B) -1
 - C) 1
 - D) 2

(EMC01-0009E)

- If α and β are the roots of $4x^2 + 3x + 7 = 0$, then the value of $\frac{1}{\alpha} + \frac{1}{\beta}$ is

- A) $-\frac{3}{7}$
- B) $-\frac{3}{4}$
- C) $\frac{3}{7}$
- D) $\frac{4}{7}$

(EMC01-0010H)

- If α and β are the roots of the equation $x^2 + 2x + 4 = 0$, then $\frac{1}{\alpha^3} + \frac{1}{\beta^3}$ is equal to

- A) $-\frac{1}{2}$
- B) $\frac{1}{2}$
- C) $\frac{1}{4}$
- D) $\frac{1}{6}$

(EMC01-0011H)

- Let $p, q \in \{1, 2, 3, 4, 5\}$. The number of equations of the form $px^2 + qx + 1 = 0$ having real roots, is

- A) 7
- B) 8
- C) 9
- D) 12

(EMC01-0012M)

- If the difference of the roots of the equation $x^2 + px + 8 = 0$ is 2, then p equals

- A) ± 2
- B) $-6, 2$
- C) $-2, 6$
- D) ± 6

BUY SUBSCRIPTION TO UNLOCK ALL THE CONTENT

0349-7869821

Page 3 of 8



ONLINE.CHAJAO.COM

(EMC01-0013M)

- If one root of the equation $x^2 + px + q = 0$ is $2 + \sqrt{3}$, then the values of p and q are

- A) $-2, -\sqrt{3}$
- B) $-4, 1$
- C) $2, \sqrt{3}$
- D) $4, -1$

(EMC01-0014M)

- For what value of m , the ratio of the roots of the equation $12x^2 - mx + 5 = 0$ is $3 : 2$?

- A) $5\sqrt{10}$
- B) $10\sqrt{5}$
- C) $25\sqrt{2}$
- D) $15\sqrt{5}$

(EMC01-0015M)

- If $(1 - p)$ is a root of the quadratic equation $x^2 + px + (1 - p) = 0$, then its roots are

- A) $-1, 2$
- B) $-1, 0$
- C) $-1, 1$
- D) $0, 1$

(EMC01-0016H)

- The roots of equation $2^{2x} - 10 \cdot 2^x + 16 = 0$ are

- A) $1, 3$
- B) $1, 8$
- C) $2, 3$
- D) $2, 8$

(EMC01-0017M)

- The solution set of $\sqrt{2x-6} + \sqrt{x+4} = 5$ is

- A) $\{5\}$
- B) $\{0, 5\}$
- C) $\{1, 3\}$
- D) $\{3, 5\}$

(EMC01-0018E)

- Remainder of $x^{64} + x^{27} + 1$ when divided by $x + 1$ is

- A) 0
- B) 1
- C) 2
- D) 3

(EMC01-0019M)

- $x^2 + x + 1 + 2k(x^2 - x - 1)$ is a perfect square for how many values of k ?

- A) 0
- B) 1
- C) 2
- D) 3

(EMC01-0020E)

- Both the roots of the equation $x^2 - x - 3 = 0$ are

- A) real and rational
- B) real and irrational
- C) real and equal
- D) imaginary roots

(EMC01-0021E)

- The value of a for which the equation $2x^2 + 2\sqrt{6}x + a = 0$ has equal roots, is

- A) $\sqrt{2}$
- B) $\sqrt{3}$
- C) 2
- D) 3

(EMC01-0022M)

- If one root of equation $x^2 + ax + 12 = 0$ is 4 while the equation $x^2 + ax + b = 0$ has equal roots, then the value of b is

- A) $\frac{4}{7}$
- B) $\frac{7}{4}$
- C) $\frac{4}{49}$
- D) $\frac{49}{4}$

(EMC01-0023H)

- One root of $mx^2 - 14x + 8 = 0$ is 6 times the other root. Then m is equal to

- A) 1
- B) 2
- C) 3
- D) None of these



(EMC01-0024M)

- If $\alpha + \beta = 4$ and $\alpha^3 + \beta^3 = 44$, then α, β are the roots of the equation

- A) $2x^2 - 7x + 6 = 0$
- B) $3x^2 - 12x + 5 = 0$
- C) $4x^2 + 22x + 15 = 0$
- D) $9x^2 - 27x + 20 = 0$

PARHO AUR
Chaa�ao

(EMC01-0025M)

- If $\{\alpha, \beta\}$ is the solution set of $2x^2 - 3x + 5 = 0$, then the equation with solution set $\{\alpha^2 + 1, \beta^2 + 1\}$ is

- A) $4x^2 - 3x - 18 = 0$
- B) $4x^2 - 3x + 18 = 0$
- C) $4x^2 - 3x + 18 = 0$
- D) $4x^2 + 3x + 18 = 0$



PARHO AUR
Chaa�ao

BUY SUBSCRIPTION TO UNLOCK ALL THE CONTENT

0349-7869821

Page 7 of 8



ONLINE.CHAAJAO.COM

Answers Key	
1	B
2	A
3	C
4	B
5	C
6	D
7	B
8	D
9	A
10	C
11	D
12	D
13	B
14	A
15	B
16	A
17	A
18	B
19	C
20	B
21	D
22	D
23	C
24	B
25	D